



January 2013 (updated May 2015)

ELDER CARE

A Resource for Interprofessional Providers

Not All Dementia is Alzheimer's Disease

Richard Caselli, MD, Department of Neurology, Mayo Clinic Scottsdale

Alzheimer's disease is a common problem that is appropriately garnering both public and professional attention. As attention to Alzheimer's increases, however, there is a risk that other causes of dementia-like illnesses will be overlooked, including the 10-15% of these cases that may be reversible. Examples of dementia-like illnesses that may be confused with Alzheimer's disease are shown in Table 1.

There are four features of dementia-like illnesses that should raise concern about causes other than Alzheimer's disease. They include (1) speed of onset, (2) age of onset, (3) the patient's cognitive and neurological profile, and (4) a medical history that might indicate a non-Alzheimer's cause.

Speed of Onset and Progression

The more rapid the onset, the less likely a dementia-like condition is Alzheimer's disease. Examples of conditions confused with Alzheimer's but which have a more rapid-onset or progression range from delirium to mercury toxicity (Table 1).

Age at Onset

The younger the age at onset, the greater the chance of a non-Alzheimer's dementia. This is particularly true if there is no family history of early-onset Alzheimer's.

Cognitive and Neurological profile

Alzheimer's disease is a slowly progressive disorder that begins by affecting memory and gradually affects other domains including executive skills and naming. Anosognosia (lack of awareness of one's disability) may also develop, as may personality changes; but, gait and non-neurological functions are usually spared. Patterns different than this should raise the possibility of a non-Alzheimer's dementia.

Medical History

A patient's medical history and medications should be considered. Many medical conditions, ranging from cerebrovascular disease to drug side effects, can cause a dementia-like illness that is confused with Alzheimer's.

Table 1. Examples of Dementia-Like Illnesses that Can Be Confused with Alzheimer's Disease

Speed of Onset			
Underlying Cause	Acute	Subacute	Chronic
Degenerative	Delirium	ALS-dementia with hypoventilation	See Table 2
Infectious	Viral encephalitis	Fungal meningitis	Neurosyphilis
Inflammation	Disseminated Encephalomyelitis	Paraneoplastic syndrome	Autoimmune encephalopathy
Neoplastic	Obstructive hydrocephalus	Glioblastoma	Orbitofrontal meningioma
Nutritional	Wernicke-Korsakoff		Vitamin B12 deficiency
Psychiatric	Acute psychosis	Inadequately controlled psychosis	Severe depression
Toxic	Drug or alcohol intoxication	Mercury toxicity	Polypharmacy
Traumatic	Acute head injury	Subdural hematoma	Chronic traumatic encephalopathy
Vascular	Acute stroke	Disseminated intravascular coagulation	Vascular dementia

TIPS FOR RECOGNIZING NON-ALZHEIMER'S DEMENTIA SYNDROMES

- When patients are being evaluated because of cognitive impairment, consider causes other than Alzheimer's disease when the impairment develops at a young age, when it develops over a short period of time, or when a movement disorder, including gait disorder, is a predominant symptom.
- Always consider potentially reversible causes of cognitive impairment, like drug side effects, infection, depression, nutritional deficiencies, and delirium.

ELDER CARE

Continued from front page

Reversible Imitators of Alzheimer's Disease

Although reversible causes account for only 10-15% of cases of dementia-like illnesses, it is essential not to overlook them. The following cases illustrate real patients who were labeled as having Alzheimer's dementia, but who subsequently turned out to have a reversible medical condition.

Case 1: Drug Toxicity A 72-year-old man with a history of bipolar disorder presented with a 7-month history of memory loss that was preceded by two years of impaired gait. He scored 18/30 on a Mini-Mental State Exam (MMSE). He was on a stable dose of lithium for years but blood work showed an elevated level. After 3 days off lithium his cognitive function and gait returned to normal.

Case 2: Infection A 66-year-old man with diabetes developed progressive memory and gait difficulties over two months. Brain imaging showed ventricular enlargement suggesting normal-pressure hydrocephalus, but the patient failed to improve after serial spinal taps. Spinal fluid cultures grew *Coccidioides immitis* that responded to fluconazole and his cognition and gait returned to normal.

Case 3. Depression A 74-year old woman developed personality changes, apathy, and difficulty with memory during the year following the death of her husband. Over the year she withdrew from social activities, and she reported difficulty participating in conversation and was sleeping poorly. Her MMSE score was 23/30, but she didn't seem to put effort into answering the questions. Her score on the Geriatric Depression Scale (GDS) was 23/30, indicating moderate-severe depression. A trial of antidepressant medication was initiated and over a period of two months, the patient's cognitive status returned to normal.

Comment All three cases were atypical for Alzheimer's disease because the onset of symptoms was too rapid. In addition, cases 1 and 2 involved a recent-onset gait disorder, which is an indicator that Alzheimer's disease is not the cause of a patient's cognitive decline. In case 3, the patient had depression - an important diagnosis to exclude because it can sometimes mimic the cognitive impairment seen in dementia.


Non-Reversible Causes of Dementia

Although Alzheimer's disease is the most common and well-known form of dementia, accounting for about 70% of cases, several other degenerative neurological disorders can cause irreversible dementia and are often confused with Alzheimer's disease. The most common of these disorders is

Table 2. Non-Reversible Dementia Syndromes That May Be Misdiagnosed as Alzheimer's Disease

Syndrome	Key Features
Vascular Dementia	<ul style="list-style-type: none">- Risk factors for cerebrovascular disease- Cerebrovascular disease on brain imaging- Psychomotor retardation w/slowed gait
Lewy Body Dementia	<ul style="list-style-type: none">- New-onset of hallucinations- Dream-enactment behaviors (e.g., yelling, kicking, spitting, flailing during REM sleep)- Parkinson-like movement disorder
Fronto-Temporal Dementia	<ul style="list-style-type: none">- Frontotemporal atrophy on imaging (see figure)- Behavioral variant: apathy, social withdrawal, disinhibition (lack of social tact)- Semantic variant (trouble finding words)

vascular dementia (now called vascular cognitive impairment), which accounts for about 15-20% of dementia cases and is often mixed with Alzheimer's. Other common syndromes include Lewy body dementia and fronto-temporal dementia, each accounting for about 10% of cases. Key characteristics of these syndromes are shown in Table 2. Although not reversible, recognizing these conditions allows for more accurate prognosis and more effective use of symptom-modifying medications.



References and Resources

- Baborie A, Griffiths TD, Jaros E, et al. Frontotemporal dementia in elderly individuals. *Archives of Neurology*. 2012; 69:1052-60.
Caselli RJ, Tariot PN. *Alzheimer's Disease and Its Variants*. New York, NY: Oxford University Press; 2010.
Potter GG, Steffens DC. Contribution of depression to cognitive impairment and dementia in older adults. *Neurologist*. 2007; 13:105-17
Zaniqni S, Calandra-Nuonaura G, Grimaldi D, Cortelli P. REM behavior disorder and neurodegenerative disease. *Sleep Medicine* 2011; 12 Suppl 2: S54-8.

Interprofessional care improves the outcomes of older adults with complex health problems

Editors: Mindy Fain, MD; Jane Mohler, NP-c, MPH, PhD; and Barry D. Weiss, MD
Interprofessional Associate Editors: Tracy Carroll, PT, CHT, MPH; David Coon, PhD; Jeannie Lee, PharmD, BCPS;
Lisa O'Neill, MPH; Floribella Redondo; Laura Vitkus, BA
The University of Arizona, PO Box 245069, Tucson, AZ 85724-5069 | (520) 626-5800 | <http://aging.medicine.arizona.edu>

Supported by: Donald W. Reynolds Foundation, Arizona Geriatric Education Center and Arizona Center on Aging

This project was supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) under grant number UB4HP19047, Arizona Geriatric Education Center. This information or content and conclusions are those of the author and should not be construed as the official position or policy of, nor should any endorsements be inferred by HRSA, HHS or the U.S. Government.