



January 2023

# ELDER CARE

## A Resource for Interprofessional Providers

### Outpatient Management of Constipation in Older Adults

Darlene Moyer, MD, HonorHealth, and Amy Tierney, MD, Scottsdale Healthcare, Scottsdale, AZ

Constipation is common in older adults. Up to 28% of individuals over age 65 experience constipation, and it is the reason for more than 2.5 million office visits to primary providers each year in the US.

Although constipation is common in older adults, it should not be considered normal. When evaluating patients who have constipation, clinicians should seek to identify reversible causes with the goal of improving quality of life and avoiding complications that include risk for fecal impaction and incontinence, hemorrhoids, anal fissure, organ prolapse, and bowel obstruction.

#### Diagnostic Criteria

Patients and clinicians often have different ideas about what is normal when it comes to bowel movements. To help define and diagnose constipation, the American Gastroenterological Association (AGA) refers to otherwise uncomplicated constipation as “functional” constipation and recommends using the Rome VI criteria for diagnosis.

The Rome IV criteria specify that the first symptoms of constipation should have begun at least 6 months previously and that two of the following must be present for at least 3 months: (a) straining, (b) lumpy or hard stools, (c) a sensation of incomplete evacuation, (d) a sensation of anorectal obstruction/blockage, (e) the need for manual maneuvers, like digital stimulation, (f) loose stools rarely present without laxative use, and (g) the patient does not meet criteria for irritable bowel syndrome (IBS). Criteria a-e should be present for at least 25% of bowel movements. A key difference between constipation and IBS is that IBS involves pain relieved by defecation, while constipation by itself is not painful.

#### History and Physical

Constipation in older adults almost always involves multiple contributing causes (Table 1), all of which should be considered during the evaluation. The history should include

questions about medical conditions, medications, prior surgeries, and pelvic floor trauma. Questions should also specifically ask about “red flags,” such as acute onset, weight loss, rectal bleeding, and a personal or family history of colorectal cancer, any of which might indicate malignancy as the cause of constipation.

**Table 1. Factors That Can Cause or Contribute to Constipation**

Medical Conditions	Medications
Anorectal pathology (fissures, strictures, prolapse, hemorrhoids)	Antacids
Cerebrovascular disease	Anticholinergics
Colon Masses/strictures	Antidepressants
Dementia	Antihistamines
Depression	Antipsychotics
Diabetes mellitus	Calcium Channel Blockers
Hypercalcemia	Calcium supplements
Hyperparathyroidism	Diuretics causing dehydration
Hypothyroidism	Iron supplements
Inflammatory bowel disease	Opiates
Multiple sclerosis	<b>Lifestyle</b>
Parkinson’s disease	Immobility
Spinal cord injury/tumors	Lack of privacy
	Low-fiber diet
	Not responding to urge to defecate
	Poor fluid intake

#### Additional Testing

The AGA recommends performing a complete blood count, thyroid function tests, and a basic metabolic panel. Colonoscopy should be performed if there is concern about cancer or if the patient is due for routine colon cancer screening. Other tests, such as colonic transit testing, anorectal manometry, and balloon expulsion testing, should only be done when patients fail a therapeutic trial of laxatives and increased dietary fiber.

#### TIPS FOR DEALING WITH CONSTIPATION IN OLDER ADULTS

- Use the Rome IV criteria to diagnose constipation.
- Identify medical conditions, medication side effects, and lifestyle issues as potentially reversible causes of constipation.
- Recommend non-pharmacological approaches including timing of bowel movements in the morning and after meals, increased physical activity, and increased dietary fiber.
- Recommend medications for treatment of constipation only if the response to non-pharmacological approaches has been inadequate.

# ELDER CARE

Continued from front page

## Non-Pharmacological Treatment

Medical conditions that may be contributing to constipation should be treated and controlled. Constipating medications should be discontinued whenever possible or changed to agents with similar action but less potential for constipation.

Counsel patients to respond to the urge to defecate when it occurs and to develop a schedule for bowel movements. The bowels are most active in the morning and 30-60 minutes after meals, so patients should be taught to take advantage of these times to use the bathroom. Institutionalized patients should be given enough time and privacy to have bowel movements. Valsalva should be avoided in cardiac patients as it can result in bradycardia and death.

Although increasing mobility is effective for preventing constipation, there is no evidence that it is effective once constipation develops. Similarly, there is no evidence to support fluid status as a factor contributing to constipation. Increasing fluid intake in older adults, many of whom have a delicate fluid balance at baseline, should be avoided. Some evidence supports use of acupressure and abdominal massage for constipation treatment.

Dietary fiber, however, is effective and intake should be increased. The increase should occur slowly (over the course of a few weeks) to the goal intake of 25-35 grams daily. Increasing too quickly can result in bloating and flatulence.

## Pharmacologic Treatment

Evidence guiding the choice of medication to treat constipation in the elderly is weak. Medications are, however, considered appropriate to use when the response to non-pharmacological treatment has been inadequate.

To decrease adverse effects, the choice of laxative should be made with the patient's comorbidities in mind. Table 2 describes some available laxatives and gives guidance on how to choose an appropriate agent. Enemas and suppositories should only be used to treat acute constipation. Patients taking laxatives should ensure good fluid intake to avoid dehydration. Those requiring opioids should receive prophylactic bowel regimens to prevent constipation. Newer prescription medications including lubiprostone, prucalopride, linaclotide, and plecanatide have not been studied well in the elderly, should not be used first-line, and should be prescribed with consideration of their potential side effects.

Evidence does not support the use of probiotics to prevent and treat constipation in the elderly, but more research is needed. Biofeedback is only effective when treating constipation caused by anorectal dysfunction and is not appropriate for patients with cognitive impairment. Surgery, such as subtotal or total colectomy for treatment of colonic inertia, should only be considered for severe recalcitrant cases.

**Table 2. Selecting Laxatives for Constipation in Older Adults**

Type of Laxative	Examples	Key Side Effects	What to Consider
Bulk Laxatives	Psyllium Methylcellulose	Bloating, flatulence, impaction above strictures	Can decrease absorption of some medications, including warfarin, aspirin, digoxin
Emollient Laxatives (Stool softeners)	Docusate Sodium Docusate Calcium	Fecal soiling	Not recommended for chronic treatment
Osmotic Laxatives	Polyethylene Glycol	Bloating, flatulence, pulmonary edema	Avoid in patients at risk for aspiration
	Lactulose, Sorbitol	Bloating, flatulence	Recommend for patients in nursing homes
	Magnesium Sulfate Magnesium Citrate	Watery stools, urgency, magnesium toxicity, hyperkalemia	Avoid in patients with renal insufficiency
Stimulant Laxatives	Bisacodyl, Senna	Cramping, gastric and rectal irritation, melanosis coli	Recommended only for short-term use; avoid in patients with bowel obstruction

## References and Resources

- Drossman D. Functional Gastrointestinal Disorders: History, Pathophysiology, Clinical Features, and Rome IV. *Gastroenterology* 2016; 150:1262-1279.
- Dukas L, Willett WC, Giovannucci EL. Association between physical activity, fiber intake, and other lifestyle variables and constipation in a study of women. *American Journal of Gastroenterology*. 2003;98:1790-6.
- Locke G, Pemberton J, Phillips S. American Gastroenterological Association Medical Position Statement: Guidelines on Constipation. *Gastroenterology*. 2000;119:1761-1778.
- Longstreth G, Thompson W, Chey W, et al. Functional Bowel Disorders. *Gastroenterology* 2006;130:1480-1491.
- Lucak S, Lunsford T, Harris L. Evaluation and Treatment of Constipation in the Geriatric Population. *Clin Geriatr Med*. 2021; 37:85-102.
- Mounsey A, Raleigh M, Wilson A. Management of Constipation in Older Adults. *Am Fam Physician*. 2015;92(6):500-504.

## Interprofessional care improves the outcomes of older adults with complex health problems.

Editors: Mindy Fain, MD; Jane Mohler, NP-c, MPH, PhD; and Barry D. Weiss, MD

Interprofessional Associate Editors: Tracy Carroll, PT, CHT, MPH; David Coon, PhD; Marilyn Gilbert, MS, CHES; Teri Kennedy, PhD, MSW, LCSW, ACSW; Jeannie Lee, PharmD, BCPS; Marisa Menchola, PhD; Francisco Moreno, MD; Lisa O'Neill, MPH; Floribella Redondo; Laura Vitkus, BA

The University of Arizona, PO Box 245069, Tucson, AZ 85724-5069 | (520) 626-5800 | <http://aging.arizona.edu>

Supported by: Donald W. Reynolds Foundation, Arizona Geriatrics Workforce Enhancement Program and the University of Arizona Center on Aging

This project was supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) under grant number U1QHP28721, Arizona Geriatrics Workforce Enhancement Program. This information or content and conclusions are those of the author and should not be construed as the official position or policy of, nor should any endorsements be inferred by HRSA, HHS or the U.S. Government.