



February 2023

ELDER CARE

A Resource for Interprofessional Providers

Sexual Health in Older Adults - Part 2

Common Concerns, Chronic Diseases, and Sexually Transmitted Infections

Travis Glenn MD; Monica Chung MD; Darlene Moyer MD; HonorHealth Family Medicine Residency, Phoenix, AZ

In the first of our two-part series on sexual health in older adults, we addressed the importance of taking a sexual history and how to effectively do so in a culturally appropriate and sensitive way. In this edition of Elder Care, we focus on common sexual health issues in older adults and how to address them - an important concern given the increasing percentage of older adults in our communities and practices.

Common Performance Concerns

Physiologic changes associated with aging can lead to sexual performance concerns for both men and women (Table 1).

In women, decreasing estrogen causes changes in the vaginal epithelium, sometimes leading to dyspareunia. Treatment can include lubricants, vaginal estrogens, toys, or vaginal dilators. If patients opt for vaginal estrogens, it is important to counsel them appropriately on systemic absorption, risks, and benefits.

In men, erectile dysfunction (ED) is the main performance concern. ED can have organic, psychogenic, or mixed causes and treatment differs depending on the cause. Common organic causes of ED include low testosterone levels or microvascular disease. Microvascular disease can be treated with phosphodiesterase inhibitors while low testosterone may respond to hormone replacement therapy (HRT). Men should be counseled, however, on the potential risks of HRT such as prostate enlargement increased risk of cardiovascular events.

Both men and women can suffer from "widower syndrome," which is a period of abstinence after the loss of a long-term partner. Another common complaint is decreased libido. Given many causes for this, it is important to take a comprehensive history to screen for depression and anxiety.

Older adults who identify as LGTBQIA+ lived during a time of significant change from civil rights movements. It is estimated that 15% of older gay men suffered from conversion therapy. Traumatized adults may not disclose their sexual orientation for fear of unequal treatment by providers or at care facilities. Including gender pronouns in routine patient and provider introductions as well as intake forms can help patients feel more safe in the office space.

Chronic Disease and Sex

Rising rates of obesity, cardiovascular disease (CVD), and diabetes combined with longer life expectancy have revealed an interesting interplay of chronic disease and sex (Table 2). As noted, microvascular disease is one of the common reasons for ED, and it is often caused by poorly controlled CVD risk factors like hypertension and diabetes. A key to preventing ED is early and

sufficient control of those risks. Discussing these issues with patients when CVD risk factors are identified may provide patients with additional "incentives" for controlling those risk factors to prevent or delay development of ED.

Furthermore, men with a history of CVD may ask if their heart is healthy enough for sex. Current studies suggest that CVD carries a relative risk of myocardial infarction of 1.5% per sexually active men per year. Appropriate counseling should take place about what to do if patients experience chest pain during intercourse.

Another barrier may be chronic pain from arthritis or other causes. For patients whose sexual activity is being limited by pain, clinicians should discuss options for adequate pain control during intercourse.

Benign prostate hyperplasia (BPH) can cause some men to experience urinary incontinence during intercourse. If this occurs, providing adequate BPH treatment can help relieve the symptoms and the associated psychological distress. Women may also experience incontinence during sexual activity. For older women, the incontinence can be stress, urge, or mixed incontinence, any of which can be combined with pelvic organ prolapse. For women experiencing these symptoms, conduct a POP-Q (see references and resources list) to assess for level of prolapse and guide therapy as needed.

It is recommended to have early and frequent conversations about sexual health with all patients, especially for those with dementia, while they still have decision-making capacity. Additionally, some memory care institutions have developed policies regarding consent to sexual activity. The references and resource list contains a link to one of the first such policies.

Sexually Transmitted Infections in the Elderly

There are increasing rates of syphilis, gonorrhea, chlamydia, and HIV in older adults, likely due to several factors. One is that as part of the aging process, the immune system has decreases in multiple immunologic cell lines, thus inhibiting the ability to fight infections. Another reason is that current condom use is estimated at only 15% overall and significantly less for older adults. This is thought to be due to today's older adults not experiencing sex education in school during their youth, plus a lack of concern about pregnancy. Undiagnosed and untreated STIs can be mistaken as normal aging. Testing should be undertaken when screening is indicated or STI is suspected, and infections should be reported and treated. Prophylactic treatment for HIV should be offered to older adults, regardless of age, if they remain at risk. The references and resources list contains a link to the CDC's treatment guidelines.

TIPS FOR HELPING OLDER ADULTS WITH THEIR SEXUAL HEALTH

- Treating and controlling chronic disease early on can improve sexual health later in life.
- Determine whether an older adult is having difficulty with desired sexual performance because of a medical condition and offer guidance on how to address the problem.
- Educate older adults about sexually transmitted infections and proper use of condoms; they may never have been taught this information before.

ELDER CARE

Continued from front page

Final Comment

Sexual health is an important aspect of well-being that is commonly under-evaluated and under-treated in older adults. It is a multifaceted issue, encompassing physical and emotional concerns as well as raising ethical questions in those with dementia. As discussed, physical and emotional factors should be addressed as

they can be improved through treatment. Clinicians can also alleviate distress regarding sexual dysfunction by raising awareness about common symptoms. Teaching our patients about these symptoms normalizes the aging process in sexual health. In engaging our patients in open dialogue and providing a safe space to discuss their sexual health, clinicians will be better able to improve their patients' quality of life and care for the whole person.

Table 1. Common Sexual Performance Concerns

Concern	Cause	Treatment
Dyspareunia	Loss of estrogen causing vaginal atrophy, stenosis, or thinning of epithelial lining	Vaginal estrogen or lubricants
Widower Syndrome	Death of long-term partner, often followed by a period of abstinence	Reassurance, behavioral therapy, and addressing misconceptions
Erectile Dysfunction 1. Organic 2. Psychogenic 3. Mixed	1. Microvascular disease, neurogenic injury, or local contributors (e.g., penile fracture) 2. Anxiety, depression, relationship concerns, or past traumas 3. And combination of factors from both groups	Medications: Phosphodiesterase-5 inhibitors Devices: vacuums or implants Therapy: behavioral or couples counseling
Decreased Libido	Usually multifactorial	Treat underlying causes (e.g., depression)

Table 2. Chronic Conditions That Commonly Cause Sexual Dysfunction

Condition	How it affects sexual health	Way to treat or mitigate
Hypertension and cardiovascular disease	Erectile dysfunction via vascular insufficiency and reduced endothelial integrity	Individual counseling based on known risk level and health conditions along with symptoms
Diabetes	Erectile dysfunction via microvascular changes similar to hypertension and cardiovascular disease	Controlling diabetes in younger men
Benign Prostatic Hyperplasia	Outflow obstruction which can lead to: erectile dysfunction, ejaculatory dysfunction, and hypoactive sexual disease	Medications: alpha-1-blockers and 5-alpha-reductase inhibitors Office procedures: prostate steaming Surgeries: TURP and stents
Dementia	Cognitive impairment does not correspond to decreased sexual desire	Most issues surround consent, but some nursing homes are addressing this concern with patients and their families
Arthritis	Pain limits mobility	Treat chronic pain and discuss as-needed pain options
Lower Urinary Tract Dysfunction (e.g., incontinence and pelvic organ prolapse)	Often causes embarrassment about intimacy, but can also lead to dyspareunia if pelvic organ prolapse is present	Providing reassurance and treating causes of incontinence Prolapse can be managed surgically or with pessaries

References and Resources

- 2021 Sexually Transmitted Diseases Treatment Guidelines. Centers for Disease Control and Prevention. <https://www.cdc.gov/std/treatment-guidelines/default.htm>
- American Urogynecologic Society. POP_Q Tool. <https://www.augs.org/patient-services/pop-q-tool/>
- Inelmen Em, et al. The importance of sexual health in the elderly: breaking down barriers and taboos. *Aging Clin Exp Res*, 2012;24(3 Suppl):31-4
- Meanley et al. Lifetime exposure to conversion therapy and psychosocial health among midlife and older adult men who have sex with men. *Gerontologist*. 2020; 60(7):1291-1302.
- Nursing Home Sexual Expression Policy. [Resident-Sexual-Expression-Policy-revised-4-2017.pdf](https://www.riverspringliving.org/resident-sexual-expression-policy-revised-4-2017.pdf) (riverspringliving.org)
- Sex After 50. American Sexual Health Association. 2019. <http://www.ashsexualhealth.org/sexual-health/sex-after-50/>
- Sexual Rights for Seniors. American Sexual Health Association. 2019. <http://www.ashsexualhealth.org/sexual-rights-for-seniors/>
- Sexuality in Later Life. National Institute for Aging. 2019. <https://www.nia.nih.gov/health/sexuality-later-life>

Interprofessional care improves the outcomes of older adults with complex health problems

Elder Care Editors: Editor-in-Chief: Barry D Weiss, MD; Deputy Editor: Mindy Fain, MD

National Editorial Board: Theodore M Johnson II, MD, MPH, Emory University; Jenny Jordan, PT, DPT, Sacred Heart Hospital, Spokane, WA; Jane Marks, RN, MS, FNGNA, Johns Hopkins University; Josette Rivera, MD, University of California San Francisco; Jean Yudin, CRNP, University of Pennsylvania

Interprofessional Associate Editors: Tracy Carroll, PT, CHT, MPH; David Coon, PhD; Carleigh High, PT, DPT; Jeannie Lee, PharmD, BCPS; Marisa Menchola, PhD; Francisco Moreno, MD; Linnea Nagel, PA-C, MPAS; Lisa O'Neill, DBH, MPH; Floribella Redondo

Published by: The University of Arizona Center on Aging, PO Box 245027, Tucson, AZ 85724-5027 | (520) 626-5800 | <http://aging.arizona.edu>