



March 2023

ELDER CARE

A Resource for Interprofessional Providers

Delirium

Rosemary Browne, MD, College of Medicine, University of Arizona

Delirium is an acute confusional state that occurs in response to physiologic stress, most commonly from a medical illness. It is a clinical syndrome seen frequently in older adults, particularly in hospital settings, presenting in up to 30% of older adult inpatients and 70% of older adults in critical care units. Delirium prolongs hospital stays, accelerates functional and cognitive decline, and increases rates of death and institutionalization. Delirium is often missed clinically. Preventing delirium is a primary focus of Age-Friendly Care for hospitalized older adults.

Risk Factors

Predisposing factors include age > 80 years, prior history of delirium, male gender, immobility, sensory impairment, complex chronic illness, polypharmacy, and malnutrition. In addition, patients with mild cognitive impairment, dementia movement disorders, cerebrovascular disease, or chronic substance abuse are also at increased risk.

Precipitating factors include conditions such as an acute illness, metabolic derangements, use of physical restraints, dehydration, pain, urinary retention and fecal impaction. Infection commonly presents as delirium, and is a poor prognostic sign in older adults with COVID-19. Surgical patients can experience delirium in the perioperative period. In addition, delirium can result from changes in medication and substance use and abuse. Medications associated with delirium are shown in Table 1.

Diagnosis

The fifth edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) defines delirium as (1) a disturbance in attention that (2) develops over a short period of time and (3) is accompanied by a disturbance in cognition. It adds that (4) these disturbances are not better explained by a pre-existing, established, or evolving neurocognitive disorder, and do not occur in the context of a severely reduced level of arousal such as coma. Also, that (5) there should also be evidence from history, physical exam, or lab testing that the disturbances are caused by a medical condition or exposure to a medication or toxin. Note that newly diagnosed dementia can sometimes be confused for delirium. Table 2 lists key differences.

Delirium can have different clinical presentations. Overt agitation and behavior disturbances make hyperactive delirium easy to diagnose. More commonly, however, older adults present in a quiet, hypoactive state, their delirium less evident on exam, making the diagnosis easy to miss. With mixed delirium, patients often fluctuate between the two. Delirium assessment tools can help to identify hyperactive, hypoactive, and mixed types.

The Confusion Assessment Method (CAM) is a widely used, evidence based instrument for delirium diagnosis. It is well-validated with high sensitivity, specificity, and inter-rater reliability. The CAM has four components. To diagnose delirium, the first two must be present: (1) acute onset of changing or rapidly fluctuating mental status, and (2) inattention, along with at least one of the next two components (3) disorganized thinking and/or (4) altered level of consciousness (Table 3).

<p>Neuropsychiatric Medications</p> <ul style="list-style-type: none"> • Anticonvulsants • Antidepressants • Antipsychotics • Dopamine agonists <p>Gastrointestinal Medications</p> <ul style="list-style-type: none"> • Antiemetics • Antispasmodics • H-2 blockers <p>Cardiovascular Medications</p> <ul style="list-style-type: none"> • Antiarrhythmics • Beta blockers • Clonidine • Digoxin • Diuretics <p>Analgesics</p> <ul style="list-style-type: none"> • Opioids • Non-steroidal anti-inflammatory medications 	<p>Allergy Medications</p> <ul style="list-style-type: none"> • Anticholinergics • Antihistamines <p>Sedatives</p> <ul style="list-style-type: none"> • Benzodiazepines and all other sedative-hypnotics <p>Herbal Medicines</p> <ul style="list-style-type: none"> • Atropa belladonna • Burdock root • Black henbane • Jimson weed • Mandrake • St. John's Wort • Valerian <p>Miscellaneous</p> <ul style="list-style-type: none"> • Corticosteroids • Hypoglycemics • Muscle relaxants • Mydriatics
--	--

TIPS FOR DEALING WITH DELIRIUM IN OLDER ADULTS

- Preventing delirium is the best intervention in the care of older adults. Multicomponent interprofessional non-pharmacologic delirium prevention strategies (Table 4) can help.
- Screen frequently for delirium in a hospitalized older adult.
- The CAM is a well-validated tool for confirming the diagnosis of delirium.
- Treatment of the underlying condition is paramount to resolving the delirium.
- Evaluation must include a medication review to identify potential contributory drugs.

ELDER CARE

Continued from front page

Characteristic	Delirium	Dementia
Onset	Acute	Insidious
Cognitive Dysfunction	Obvious	Can be subtle in early dementia
Mental Status	Fluctuating	Progressive impairment
Reversibility	Potentially Reversible	Irreversible

Treatment

Inter-professional **prevention** approaches (Table 4) are the most important interventions for delirium. Once delirium is diagnosed, treatment should be instituted immediately, as the duration of delirium is associated with increasing risk of poor outcomes.

Resolving underlying medical problems is paramount for treating delirium. Patients should be evaluated for treatable causes (see precipitating factors listed previously). Always review medications, especially any recently started. Further testing, including brain imaging, lumbar puncture, and/or EEG should follow if a treatable cause is not identified. Even after treating a causal condition, however, resolution of delirium may take weeks to months to resolve, and sometimes longer.

Behavioral interventions (Table 4) can be effective not only for preventing delirium, but also for treating delirium once it has developed. Physical restraints can worsen symptoms and should be avoided.

Pharmacologic therapy is often used in the treatment of delirium, but its role and value vary with clinical situations. Medication may reduce agitation, but prolong the delirium and cognitive decline. Hypoactive delirium is managed by treating reversible medical conditions and instituting the interventions listed in Table 4.

In contrast, hyperactive delirium can require emergent intervention to prevent harm. Although no medications are FDA approved for treating delirium, anti-psychotics are used when no other measures are successful. While haloperidol has long been considered the drug of choice for

References and Resources

- Andersen-Ranberg NC, Poulsen LM, Perner A, et al. Haloperidol for the treatment of delirium in ICU patients. *NEJM*. 2022 Dec 29;387(26):2425-35. Confusion Assessment Method Training Manual. https://americandeliriumsociety.org/wp-content/uploads/2021/08/CAM-Long_Training-Manual.pdf
- Marcantonio, E.R. Delirium in hospitalized older patients. *NEJM*. 2017;337: 1456-66.
- Mattison, M. In the Clinic: Delirium. *Ann. Int. Med.* 2020; 173 (7): ITC49-ITC64. <https://doi.org/10.7326/AITC202010060>
- Oh, ES, Fong, TG, Hsieh, TT, Inouye, SK. Delirium in older persons-advances in diagnosis and treatment. *JAMA*. 2017; 318: 1161-74.
- Tyson B, Shahein A, Erdodi L, et al. Delirium as a Presenting Symptom of COVID-19. *Cogn Behav Neurol*. 2022 Jun 1;35(2):123-129. doi: 10.1097/WNN.0000000000000305. PMID: 35486533; PMCID: PMC9148631.
- Wilson, JE, Mart, MF, Cunningham, C et al. Delirium. *Nat Rev Dis Primers*. 2020; 6 (90). <https://doi.org/10.1038/s41572-020-00223-4>

hyperactive delirium and it can be administered by multiple routes (PO, IM, IV), recent data suggest that the newer antipsychotics (e.g., quetiapine risperidone, olanzapine) result in more rapid symptom improvement and are better tolerated by patients, including fewer side effects. Benzodiazepines are indicated only for the treatment of delirium from alcohol or benzodiazepine withdrawal.

Table 3. How to Assess the Diagnostic Criteria in the Confusion Assessment Method (CAM)

- **Acute Change or Fluctuation in Mental Status:** Assess by history and observation. Staff and family can attest to the admission/pre-op or pre-hospital cognitive status of the patient. Any acute confusional state should make the provider consider delirium.
- **Inattention:** Is the patient able to answer a direct question with an appropriate answer? Can the patient stay “on track” in normal conversation? If the answer is no, also look for fluctuations in levels of attention, which can further signal delirium.
- **Disorganized Thinking:** Is the patient’s speech/thought process rambling, unclear, unpredictable, illogical, and/or irrelevant?
- **Altered Level of Consciousness:** Assess the patient for alertness, vigilance, lethargy, stupor, or coma.

Table 4. Prevention and Treatment of Delirium

- Orientation: provide clock, calendar, white board, staff re-orientation
- Cognitive stimulation: provide familiar visitors, activities
- Improve sleep: avoid nighttime vitals and blood draws, provide quiet environment
- Early mobilization in hospital: reduce immobility, use occupational and physical therapy
- Nutrition and hydration (don’t forget dentures)
- Avoid restraints, tethers, urinary catheters
- Facilitate communication with hearing aids, spectacles
- Avoid medications with anticholinergic effects
- Manage pain; do not over- or under-treat
- Assess for new medical issues and treat immediately

Interprofessional care improves the outcomes of older adults with complex health problems.

Editors: Mindy Fain, MD; Jane Mohler, NP-c, MPH, PhD; and Barry D. Weiss, MD

Interprofessional Associate Editors: Tracy Carroll, PT, CHT, MPH; David Coon, PhD; Marilyn Gilbert, MS, CHES;

Jeannie Lee, PharmD, BCPS; Marisa Menchola, PhD; Francisco Moreno, MD; Linnea Nagel, PA-C, MPAS; Lisa O’Neill, DBH, MPH; Floribella Redondo; Laura Vitkus, MPH

The University of Arizona, PO Box 245069, Tucson, AZ 85724-5069 | (520) 626-5800 | <http://aging.arizona.edu>

Supported by: Donald W. Reynolds Foundation, Arizona Geriatrics Workforce Enhancement Program and the University of Arizona Center on Aging

This project was supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) under grant number U1QHP28721, Arizona Geriatrics Workforce Enhancement Program. This information or content and conclusions are those of the author and should not be construed as the official position or policy of, nor should any endorsements be inferred by HRSA, HHS or the U.S. Government.